

Letter to the Editor

***Pantoea* clinical isolates cannot be accurately assigned to species based on metabolic profiling**

F. Rezzonico, V.O. Stockwell, M. Tonolla, B. Duffy, T.H.M. Smits.
Pantoea clinical isolates cannot be accurately assigned to species based on metabolic profiling.
Transpl Infect Dis 2011. All rights reserved

**F. Rezzonico¹, V.O. Stockwell²,
M. Tonolla³, B. Duffy¹, T.H.M. Smits¹**

¹Plant Protection Division, Agroscope Changins-Wädenswil ACW, Wädenswil, Switzerland, ²Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon, USA, ³Microbial Ecology Group, Microbiology Unit, BIVEG Department UNIGE, Cantonal Institute of Microbiology, Bellinzona, Switzerland

Correspondence to:
Fabio Rezzonico
Agroscope Changins-Wädenswil ACW, Schloss 1
Postfach,
CH-8820 Wädenswil, Switzerland
Tel: +41 44 783 6331
Fax: +41 44 780 6341
E-mail: fabio.rezzonico@acw.admin.ch

Received 24 May 2011, accepted for publication 13 August 2011

DOI: 10.1111/j.1399-3062.2011.00684.x
Transpl Infect Dis 2011; **0**: 1–2

To the Editor

We wish to comment on a recent clinical report of *Pantoea agglomerans* (1), motivated by the enduring influence that such reports have on regulatory assessment of *P. agglomerans* biocontrol strains for beneficial agricultural application as natural plant disease control alternatives.

Pantoea and *P. agglomerans* are ecologically diverse taxa that have undergone substantial taxonomic revisions, which can confuse identification efforts (2, 3). Identification in clinical reports (1, 4) is often based on metabolic phenotyping, an inadequate approach for *Pantoea* species resolution (5, 6). Databases for standard metabolic tests such as an API system (BioMérieux SA, Marcy l'Etoile, France) lack reference depth essential to discriminate *P. agglomerans* from the other 13 *Pantoea* species, and best-available matching can deliver false identity as *P. agglomerans*. Use of API Coryne (a gram-positive Actinobacteria kit) in this report (1) probably further complicates identification.

Recently, multilocus sequencing and proteomic analyses of clinical isolates initially reported as *P. agglomerans* found that many were misidentified and deserved reassignment to other *Pantoea* species or even other enterobacterial genera (2, 5, 6).

We appreciate the impracticality of integrating additional steps in routine clinical diagnostics, but in published case reports application of the most sensitive technology must be warranted to ensure unambiguous identification. Moreover, although conservation of published isolates by authors or culture collections is imperative for independent identity confirmation as technology and taxonomy advance, obtaining even recently published clinical isolates unfortunately is rarely successful. This is a major obstacle to investigating “pathogenic potential of *P. agglomerans*” (1, 4), a recurrent inference despite historical misidentification of this species and absence of evidential support for animal pathogenicity/toxicity (2).

We hope this comment will foster dialog and collaboration between clinical and environmental bacteriology communities essential to resolve the enigma of *P. agglomerans*.

References

1. Shubov A, Jagannathan P, Chin-Hong PV. *Pantoea agglomerans* pneumonia in a heart-lung transplant recipient: case report and a review of an emerging pathogen in immunocompromised hosts. *Transpl Infect Dis* 2011; 13 (5) [Epub 2011 April 19].
2. Rezzonico F, Smits THM, Montesinos E, Frey JE, Duffy B. Genotypic comparison of *Pantoea agglomerans* plant and clinical strains. *BMC Microbiol* 2009; 9: 204.
3. Brady CL, Cleenwerck I, Venter SN, Engelbeen K, de Vos P, Coutinho TA. Emended description of the genus *Pantoea* and description of four novel species from human clinical samples, *Pantoea septica* sp. nov., *Pantoea eucrina* sp. nov., *Pantoea brenneri* sp. nov. and *Pantoea conspicua* sp. nov., and transfer of *Pectobacterium cypripedii* (Hori 1911) Brenner et al. 1973 emend. Hauben et al. 1998 to the genus *Pantoea* emend. as *Pantoea cypripedii* comb. nov. *Int J Syst Evol Microbiol* 2010; 60 (10): 2430–2440.
4. Cruz AT, Cazacu AC, Allen CH. *Pantoea agglomerans*, a plant pathogen causing human disease. *J Clin Microbiol* 2007; 45 (6): 1989–1992.
5. Rezzonico F, Vogel G, Duffy B, Tonolla M. Application of whole-cell matrix-assisted laser desorption ionization–time of flight mass spectrometry for rapid identification and clustering analysis of *Pantoea* species. *Appl Environ Microbiol* 2010; 76 (13): 4497–4509.
6. Delétoile A, Decré D, Courant S, et al. Phylogeny and identification of *Pantoea* species and typing of *Pantoea agglomerans* strains by multilocus gene sequencing. *J Clin Microbiol* 2009; 47 (2): 300–310.